



SPECIAL INSTRUCTIONS

Please follow all instructions on this sheet carefully, failure to do so could result in a cancellation or postponing procedure.

No using tobacco or marijuana after midnight prior to procedure!

DIABETIC PATIENTS

If you take insulin (pills or injections) you will take a 1/2 dose the day before the procedure and then you will NOT take any the morning of.

BLOOD THINNERS

If you are taking any blood thinners (I.e. Eliquis, Warfarin, Plavix, etc) please hold those prior to the procedure, our office will let you know how many days to hold.

TRANSPORTATION

You will not be able to drive that day, anyone 18 or older will be fine.

Public transportation is not allowed, (i.e. lyft, uber, bus, taxi, or ride-share). Good Shephard (479-283-6285) and Medicaid transportation services are allowed.

GETTING STARTED

Do not consume any fiber, nuts, corn, popcorn or salad 2-3 days before your procedure. These foods could cause your bowels to not completely clear out and could interfere with the visualization of your colon.

Stop oral iron, Metamucil, and any other fiber supplements for 3 days before the procedure.

If you are taking any type 2 diabetic medications for weight loss (i.e. Mounjaro, Ozempic, Wegovy or Trulicity) please hold for one week prior to procedure.

The day before the procedure

As soon as you wake up, you will be on a clear liquid diet until after you have had your procedure. **NO SOLID FOOD, DIARY PRODUCTS, OR RED DYE!!**

Clear liquids that are okay to have would be things like broth, strained soups, water (flavored is okay), Pedialyte, Gatorade, Powerade, Ginger Ale, Sprite, 7up, apple juice, orange juice (no pulp), coffee, tea, lemonade, jell-o, popsicles, icee, slushies, etc.

At 6:00 pm, you will begin the prep by mixing one pouch with water into container and mix. After finishing the container, you will refill it 2 more times and drink within the next hour. Chilling the prep and using a straw helps. If nauseous, stop for 1 hour and the resume drinking slowly.

The day of the procedure

You will continue the clear liquid diet as the day before. You may have liquids up to **TWO** hours prior to your arrival time.

At 4:30 am, you will take the second dose of Suflave. Repeat the same steps from the night before.

Please take your heart, seizure, and/or blood pressure medication with a small sip of water.

The procedure will **NOT** be at the GI Alliance office in Springdale. Please go to North Hills Gastroenterology Endoscopy Center in Fayetteville, their address is 3344 N. Futrall Dr. Suite 3. **Please arrival 45 minutes early.**

DATE:

TIME:

PHYSICIAN: